

Attorney Docket No. 05725.1225-00000

Customer No.



22852

PATENT TRADEMARK OFFICE

UNITED STATES PATENT APPLICATION

OF

STÉPHANE SABELLE, LAURE RAMOS, AND MADELEINE LEDUC

FOR

**PARA-PHENYLENEDIAMINE DERIVATIVES CONTAINING A PYRROLIDYL GROUP,
AND USE OF THESE DERIVATIVES FOR COLORING KERATIN FIBERS**

[001] This application claims benefit of U.S. Provisional Application No. 60/408,900, filed September 9, 2002.

[002] Disclosed herein are novel pyrrolidyl-substituted para-phenylenediamine derivatives. Also disclosed herein are dye compositions comprising the novel pyrrolidyl-substituted para-phenylenediamine derivatives and a process for dyeing keratin fibers using the dye compositions.

[003] It is known practice to dye keratin fibers, for example, human hair, with dye compositions comprising oxidation dye precursors, for example, oxidation dye precursors chosen from ortho- and para-phenylenediamines; ortho- and para-aminophenols; and heterocyclic compounds (such as diamino-pyrazole derivatives, pyrazolo[1,5-a]pyrimidine derivatives, pyrimidine derivatives, pyridine derivatives, 5,6-dihydroxyindole derivatives and 5,6-dihydroxyindoline derivatives). Oxidation dye precursors, or oxidation bases, are colorless or weakly colored compounds which, when combined with oxidizing products, can give rise to colored compounds and dyes by a process of oxidative condensation.

[004] It is also known that the shades obtained with oxidation bases may be varied by combining the oxidation bases with couplers or coloration modifiers. For example, coloration modifiers may be chosen from meta-phenylenediamines, meta-aminophenols, meta-hydroxyphenols and certain heterocyclic compounds, for instance pyrazolo[1,5-b]-1,2,4-triazole derivatives, pyrazolo[3,2-c]-1,2,4-triazole derivatives, pyrazolo[1,5-a]pyrimidine derivatives, pyridine derivatives, 5-pyrazolone derivatives, indoline derivatives and indole derivatives.

[005] The variety of molecules that can be used as oxidation bases and couplers allows a wide range of colors to be obtained.

[006] The "permanent" coloration obtained using these oxidation dyes must satisfy a certain number of requirements. For example, the oxidation dyes should not have toxicological drawbacks. In addition, oxidation dyes should produce shades in the desired intensity, and the shades should show good resistance to external agents (for instance, light, bad weather, washing, permanent-waving, perspiration and rubbing).

Permanent oxidation dyes should also be able to cover grey hair and, should be as unselective as possible, i.e., ideally they should produce the smallest possible differences in coloration along the same length of a keratin fiber, which may in fact be differently sensitized (i.e., damaged) between its end and its root. It is also desired that oxidation dyes show good chemical stability in the formulations and have a good toxicological profile.

[007] In the field of hair dyeing, para-phenylenediamine and para-tolylenediamine are oxidation bases that are widely used. They can produce varied shades with oxidation couplers.

[008] However, there is a need to discover novel oxidation bases with a better toxicological profile than that of para-phenylenediamine and para-tolylenediamine, while at the same time being able to give the keratin fibers excellent properties in terms of at least one of: color intensity, variety of shades, color uniformity and resistance to external agents.

[009] It is already known to use para-phenylenediamine derivatives substituted with a pyrrolidine group as oxidation bases for coloring keratin fibers. For example, U.S. Patent No. 5,851,237 describes the use of 1-(4-aminophenyl)pyrrolidine derivatives optionally substituted on the benzene nucleus as replacements for para-phenylenediamine. U.S. Patent No. 5,993,491 proposes the use of N-(4-aminophenyl)-2-hydroxymethylpyrrolidine derivatives optionally substituted on the benzene nucleus and on the pyrrolidine heterocycle in position 4 with a hydroxyl radical, as replacements for para-


phenylenediamine. Patent application JP 11-158 048 proposes compositions containing at least one compound chosen from 4-aminoaniline derivatives optionally substituted on the benzene nucleus and one of the nitrogen atoms of which is included in a 5- to 7-membered carbon-based ring. Patent application EP 1 200 052 describes dye compositions containing, as an oxidation base, para-phenylenediamine derivatives in which one of the amino groups forms a pyrrolidine ring substituted in position 3 with an amino group.

[010] However, the previously disclosed compounds in the art may not give the hair a coloration equivalent in quality to that obtained with para-phenylenediamine or with para-tolylenediamine due to a lack of intensity and lack of uniformity of the color.

[011] There is thus a real need for novel oxidation bases that simultaneously have a good toxicological profile and properties such that the compositions comprising the novel oxidation bases can give the hair excellent properties in terms of at least one of: color intensity, variety of shades, color uniformity and resistance with respect to various external attacking factors to which the hair may be subjected.

[012] Thus, the present disclosure describes novel dye compositions that do not have the drawbacks of the oxidation bases of the prior art. These novel dye compositions for dyeing keratin fibers do not degrade the keratin fibers, while at the same time are capable of producing intense colorations in varied shades, which are relatively unselective, particularly resistant and show a good toxicological profile.

[013] One embodiment of the invention is a compound chosen from para-phenylenediamine derivatives substituted with a pyrrolidyl group, wherein said pyrrolidyl-substituted para-phenylenediamine derivative is chosen from derivatives corresponding to formula (I), and the addition salts thereof,



wherein:

- n is an integer ranging from 0 to 4, provided that when n is greater than or equal to 2, then the radicals R_1 may be identical or different,
- R_1 is chosen from halogen atoms; saturated and unsaturated aliphatic and alicyclic C_1 - C_6 hydrocarbon-based chains, wherein at least one carbon atom of the hydrocarbon-based chain may be replaced with at least one entity chosen from oxygen, nitrogen, silicon and sulphur atoms, and from SO and SO_2 groups, provided that the radical R_1 does not comprise a peroxide bond or a diazo, nitro or nitroso radical; and further wherein the hydrocarbon-based chain may be substituted with at least one group chosen from halogen atoms, hydroxyl, amino, mono-, di(C_1 - C_4)alkylamino and tri(C_1 - C_4)alkylammonium radicals;
- Y is chosen from a covalent bond and a C_1 - C_{14} alkylene chain that may be linear or branched, wherein at least one carbon atom of the alkylene chain may be replaced with at least one atom chosen from oxygen, nitrogen and sulphur atoms, or with a group chosen from SO and SO_2 groups; wherein the alkylene chain may be substituted with at least one radical chosen from hydroxyl, C_1 - C_6 alkoxy, amino, C_1 - C_6 alkylamino and C_1 - C_6 dialkylamino radicals; and further wherein the chain may bear at least one ketone

functional group,

- R_2 is chosen from 3- to 7-membered saturated and unsaturated carbocycles and heterocycles, which may be substituted with at least one radical chosen from C_1 - C_6 alkyl and C_1 - C_6 hydroxyalkyl radicals.

[014] Another embodiment disclosed herein is a dye composition comprising at least one pyrrolidyl-substituted para-phenylenediamine derivative chosen from the derivatives corresponding to formula (I) as an oxidation base.

[015] Yet another embodiment disclosed herein is the use of the novel para-phenylenediamine derivatives to dye keratin fibers, as well as a process for dyeing keratin fibers, for instance, human keratin fibers such as the hair, using the composition comprising at least one pyrrolidyl-substituted para-phenylenediamine derivative chosen from derivatives corresponding to formula (I).

[016] The dye composition comprising at least one pyrrolidyl-substituted para-phenylenediamine derivative chosen from derivatives corresponding to formula (I) as disclosed herein can produce a chromatic, powerful, relatively unselective and resistant coloration of keratin fibers, and has a good toxicological profile.

[017] In the context of the disclosure herein, an aliphatic hydrocarbon-based chain can be either a linear or branched chain that may contain unsaturations of the alkene or alkyne type. An alicyclic hydrocarbon-based chain is a branched chain containing a cyclic structure, which may contain one or more unsaturations of the alkene or alkyne type, but which does not contain an aromatic ring structure. When the carbon atoms of the chain are replaced with an entity, "T" chosen from oxygen, sulphur, nitrogen and silicon atoms, and from SO and SO₂ groups, a unit -T-CH₂- or -T- may be obtained.

[018] By way of example, in one embodiment of the compound disclosed herein, R_1 may be chosen from chlorine atoms, bromine atoms, methyl, ethyl, isopropyl, vinyl, allyl, methoxymethyl, hydroxyethyl, 1-carboxymethyl, 1-aminomethyl, 2-carboxyethyl, 2-hydroxyethyl, 3-hydroxypropyl, 1,2-dihydroxyethyl, 1-hydroxy-2-aminoethyl, 1-amino-2-hydroxyethyl, 1,2-diaminoethyl, methoxy, ethoxy, allyloxy and 2-hydroxyethyloxy radicals.

[019] In another embodiment of the compound disclosed herein, n is equal to 0 or 1.

[020] According to yet another embodiment, R_1 is chosen from halogen atoms, C_1 - C_4 alkyl radicals, C_1 - C_4 hydroxyalkyl radicals, C_1 - C_4 aminoalkyl radicals, C_1 - C_4 alkoxy radicals and C_1 - C_4 hydroxyalkoxy radicals. By way of example, R_1 may be chosen from methyl, hydroxymethyl, 2-hydroxyethyl, 1,2-dihydroxyethyl, methoxy, isopropoxy and 2-hydroxyethoxy radicals.





[021] In yet another embodiment of the compound disclosed herein, when Y is a covalent bond, then R_2 is directly linked to the pyrrolidine ring. In still another embodiment, when Y is an alkylene chain, then for example, Y may be chosen from C_1 - C_8 alkylene chains that may comprise one or more units chosen from $-O-$, $-NR'-$ and $-NR'CO-$, wherein R' is chosen from hydrogen atoms or C_1 - C_4 alkyl radicals. Y may also be an entity chosen from: $-O-$, $-NR'-$, $-S-$, $-SO-$, and $-SO_2$.











[022] By way of another example, the radical R_2 may be a ring chosen from imidazole, pyrrolidine, piperazine, piperidine, triazole, diazepan, pyrazole, morpholine, tetrazole, furan, thiophene, pyrrole, oxazole, thiazole, and all the corresponding rings obtained by reducing at least one double bond of the rings thereof. As a further example, the triazole rings may be chosen from 1,2,3-triazole and 1,2,4-triazole rings.









[023] According to another embodiment of the compound disclosed herein, R_2 is a nitrogenous heterocycle. For example, R_2 may be chosen from imidazole, pyrrolidine, piperazine, piperidine, triazole, diazepan and pyrazole rings. According to still another embodiment, R_2 may be linked to Y via one of the nitrogen atoms of the nitrogenous heterocycle. In an additional embodiment, when R_2 is a carbocycle, R_2 may be chosen from C_4 - C_7 cycloalkyls and aryl radicals, for example, phenyl.

[024] The compounds chosen from derivatives corresponding to formula (I) may be in the form of acid salts with strong mineral acids, for instance HCl, HBr and H_2SO_4 , or with organic acids, for instance acetic acid, lactic acid, tartaric acid, citric acid and succinic acid.

[025] Examples of further embodiments of derivatives of formula (I) that may be mentioned include:

	4-(3-Imidazol-1-ylpyrrolid-1-yl)phenylamine		4-(3-[1,2,4]Triazol-1-ylpyrrolid-1-yl)phenylamine
	4-[1,3']Bipyrrolidyl-1'-yl)phenylamine		4-[3-(4-Methylpiperazin-1-yl)pyrrolid-1-yl]phenylamine

	4-[3-(4-Methyl- [1,4]diazepan-1- yl)pyrrolid-1- yl]phenylamine		4-[3-(Piperid-1- yl)pyrrolid-1- yl]phenylamine
	4-{3-[2-(4-Methyl[1,4]- diazepan-1-yl)- ethoxy]pyrrolid-1- yl}phenylamine		4-{3-[2-(4- Methylpiperazin-1- yl)ethoxy]pyrrolid-1- yl}phenylamine
	4-[3-(2-Pyrrolid-1- ylethoxy)pyrrolid-1- yl]phenylamine		4-[3-(2-Piperid-1- ylethoxy)pyrrolid-1- yl]phenylamine
	N-[1-(4-Amino- phenyl)pyrrolid-3-yl]- 2-imidazol-1- ylacetamide		N-[1-(4-Amino-3- methylphenyl)- pyrrolid-3-yl]-2- imidazol-1-yl- acetamide
	4-(3-Morpholin-4- ylpyrrolid-1- yl)phenylamine		2-Methyl-4-(3- morpholin-4-yl- pyrrolid-1-yl)- phenylamine

	[1-(4-Amino-phenyl)pyrrolid-3-yl]-(3-imidazol-1-ylpropyl)amine		2-Methyl-[1-(4-aminophenyl)-pyrrolid-3-yl]-(3-imidazol-1-ylpropyl)amine
	4-(3-Imidazol-1-ylpyrrolid-1-yl)-2-methylphenylamine		2-Methyl-4-(3-[1,2,4]triazol-1-ylpyrrolid-1-yl)phenylamine
	4-[1,3']Bipyrrolidyl-1'-yl-2-methylphenylamine		2-Methyl-4-[3-(4-methylpiperazin-1-yl)pyrrolid-1-yl]phenylamine
	2-Methyl-4-[3-(4-methyl[1,4]diazepan-1-yl)pyrrolid-1-yl]phenylamine		2-Methyl-4-[3-(piperidin-1-yl)-pyrrolid-1-yl]-phenylamine

	2-Methyl-4-{3-[2-(4-methyl[1,4]diazepan-1-yl)ethoxy]pyrrolid-1-yl}phenylamine		2-Methyl-4-{3-[2-(4-methylpiperazin-1-yl)ethoxy]pyrrolid-1-yl}phenylamine
	4-(3-Pyrazol-1-ylpyrrolid-1-yl)phenylamine		2-Methyl-4-(3-pyrazol-1-ylpyrrolid-1-yl)phenylamine
	2-Methyl-4-[3-(2-pyrrolid-1-yl)pyrrolid-1-yl]phenylamine		2-Methyl-4-[3-(2-piperid-1-ylethoxy)pyrrolid-1-yl]phenylamine
	[1-(4-Aminophenyl)pyrrolid-3-yl]cyclopentylamine		[1-(4-Aminophenyl)pyrrolid-3-yl]phenylethylamine

[026] Among these compounds, the following compounds may be mentioned:

- 4-(3-Imidazol-1-ylpyrrolid-1-yl)phenylamine
- 4-[1,3']Bipyrrolid-1'-ylphenylamine
- 4-[3-(4-Methylpiperazin-1-yl)pyrrolid-1-yl]phenylamine

- 4-{3-[2-(4-Methylpiperazin-1-yl)ethoxy]pyrrolid-1-yl}phenylamine
- 4-[3-(2-Pyrrolid-1-ylethoxy)pyrrolid-1-yl]phenylamine
- 4-[3-(Piperidin-1-yl)pyrrolid-1-yl]phenylamine
- 4-[3-(2-Piperid-1-ylethoxy)pyrrolid-1-yl]phenylamine
- [1-(4-Aminophenyl)pyrrolid-3-yl](3-imidazol-1-ylpropyl)amine
- 2-Methyl-[1-(4-aminophenyl)pyrrolid-3-yl]-(3-imidazol-1-ylpropyl)amine
- 4-(3-Imidazol-1-ylpyrrolid-1-yl)-2-methylphenylamine
- 4-[1,3']Bipyrrolid-1'-yl-2-methylphenylamine
- 2-Methyl-4-[3-(4-methylpiperazin-1-yl)pyrrolid-1-yl]phenylamine
- 2-Methyl-4-[3-(piperidin-1-yl)pyrrolid-1-yl]phenylamine
- 2-Methyl-4-{3-[2-(4-methylpiperazin-1-yl)ethoxy]pyrrolid-1-yl}phenylamine
- 4-(3-Pyrazol-1-ylpyrrolid-1-yl)phenylamine
- 2-Methyl-4-(3-pyrazol-1-ylpyrrolid-1-yl)phenylamine
- 2-Methyl-4-[3-(2-pyrrolid-1-ylethoxy)pyrrolid-1-yl]phenylamine
- 2-Methyl-4-[3-(2-piperid-1-ylethoxy)pyrrolid-1-yl]phenylamine.

[027] The dye composition as disclosed herein comprises, in a cosmetically acceptable medium that is suitable for dyeing keratin fibers, for example, human keratin fibers, at least one oxidation base comprising at least one pyrrolidyl- substituted para-phenylenediamine derivative chosen from derivatives corresponding to formula (I) as defined above. .

[028] The at least one oxidation base as disclosed above can be present in the dye composition an amount approximately ranging from 0.001% to 10%, by weight of each oxidation base, relative to the total weight of the dye composition. For example, the at least

one oxidation base may be present in an amount approximately ranging from 0.005% to 6% by weight of each oxidation base, relative to the total weight of the dye composition.

[029] The dye composition disclosed herein may also comprise at least one coupler conventionally used for dyeing keratin fibers. For example, the at least one coupler may be chosen from meta-phenylenediamines, meta-aminophenols, meta-diphenols, naphthalene-based couplers and heterocyclic couplers, and the addition salts thereof.

[030] Further examples of couplers that may be mentioned include 2-methyl-5-aminophenol, 5-N-(β -hydroxyethyl)amino-2-methylphenol, 6-chloro-2-methyl-5-aminophenol, 3-aminophenol, 1,3-dihydroxybenzene, 1,3-dihydroxy-2-methylbenzene, 4-chloro-1,3-dihydroxybenzene, 2,4-diamino-1-(β -hydroxyethyloxy)benzene, 2-amino-4-(β -hydroxyethylamino)-1-methoxybenzene, 1,3-diaminobenzene, 1,3-bis(2,4-diaminophenoxy)propane, 3-ureidoaniline, 3-ureido-1-dimethylaminobenzene, sesamol, 1- β -hydroxyethylamino-3,4-methylenedioxybenzene, α -naphthol, 2-methyl-1-naphthol, 6-hydroxyindole, 4-hydroxyindole, 4-hydroxy-N-methylindole, 2-amino-3-hydroxypyridine, 6-hydroxybenzomorpholine, 3,5-diamino-2,6-dimethoxypyridine, 1-N-(β -hydroxyethyl)-amino-3,4-methylenedioxybenzene and 2,6-bis(β -hydroxyethylamino)toluene and the acid addition salts thereof.

[031] In the dye composition disclosed herein, the optional at least one coupler may be present in an amount ranging approximately from 0.001% to 10% by weight, relative to the total weight of the dye composition. For example the at least one coupler may be present in an amount ranging approximately from 0.005% to 6% by weight, relative to the total weight of the dye composition.

[032] The dye composition disclosed herein may also comprise at least one additional oxidation base conventionally used in oxidation dyeing. By way of example,

these additional oxidation bases can be chosen from para-phenylenediamines other than those described above, bis(phenyl)alkylenediamines, para-aminophenols, bis-para-aminophenols, ortho-aminophenols, heterocyclic bases, and the addition salts thereof.

[033] For further example, mention may be made of para-phenylenediamines chosen from: para-phenylenediamine, para-tolylenediamine, 2-chloro-para-phenylenediamine, 2,3-dimethyl-para-phenylenediamine, 2,6-dimethyl-para-phenylenediamine, 2,6-diethyl-para-phenylenediamine, 2,5-dimethyl-para-phenylenediamine, N,N-dimethyl-para-phenylenediamine, N,N-diethyl-para-phenylenediamine, N,N-dipropyl-para-phenylenediamine, 4-amino-N,N-diethyl-3-methylaniline, N,N-bis(β -hydroxyethyl)-para-phenylenediamine, 4-N,N-bis(β -hydroxyethyl)amino-2-methylaniline, 4-N,N-bis(β -hydroxyethyl)amino-2-chloroaniline, 2- β -hydroxyethyl-para-phenylenediamine, 2-fluoro-para-phenylenediamine, 2-isopropyl-para-phenylenediamine, N-(β -hydroxypropyl)-para-phenylenediamine, 2-hydroxymethyl-para-phenylenediamine, N,N-dimethyl-3-methyl-para-phenylenediamine, N-ethyl-N-(β -hydroxyethyl)-para-phenylenediamine, N-(β , γ -dihydroxypropyl)-para-phenylenediamine, N-(4'-aminophenyl)-para-phenylenediamine, N-phenyl-para-phenylenediamine, 2- β -hydroxyethyloxy-para-phenylenediamine, 2- β -acetaminoethyloxy-para-phenylenediamine, N-(β -methoxyethyl)-para-phenylenediamine, 4-aminophenylpyrrolidine, 2-thienyl-para-phenylenediamine, 2- β -hydroxyethylamino-5-aminotoluene and 3-hydroxy-1-(4'-aminophenyl)pyrrolidine, and the acid addition salts thereof.

[034] Among the para-phenylenediamines mentioned above, further mention may be made of para-phenylenediamine, para-tolylenediamine, 2-isopropyl-para-phenylenediamine, 2- β -hydroxyethyl-para-phenylenediamine, 2- β -hydroxyethyloxy-para-phenylenediamine, 2,6-dimethyl-para-phenylenediamine, 2,6-diethyl-para-

phenylenediamine, 2,3-dimethyl-para-phenylenediamine, N,N-bis(β -hydroxyethyl)-para-phenylenediamine, 2-chloro-para-phenylenediamine and 2- β -acetylaminoethyloxy-para-phenylenediamine, and the acid addition salts thereof.

[035] Examples of bis(phenyl)alkylenediamines that may be mentioned, are N,N'-bis(β -hydroxyethyl)-N,N'-bis(4'-aminophenyl)-1,3-diaminopropanol, N,N'-bis(β -hydroxyethyl)-N,N'-bis(4'-aminophenyl)ethylenediamine, N,N'-bis(4-aminophenyl)tetramethylenediamine, N,N'-bis(β -hydroxyethyl)-N,N'-bis(4-aminophenyl)tetramethylenediamine, N,N'-bis(4-methylaminophenyl)tetramethylenediamine, N,N'-bis(ethyl)-N,N'-bis(4'-amino-3'-methylphenyl)ethylenediamine, 1,8-bis(2,5-diaminophenoxy)-3,6-dioxaoctane, and the acid addition salts thereof.

[036] Examples of para-aminophenols that may be mentioned, are para-aminophenol, 4-amino-3-methylphenol, 4-amino-3-fluorophenol, 4-amino-3-hydroxymethylphenol, 4-amino-2-methylphenol, 4-amino-2-hydroxymethylphenol, 4-amino-2-methoxymethylphenol, 4-amino-2-aminomethylphenol, 4-amino-2-(β -hydroxyethylaminomethyl)phenol, 4-amino-2-fluorophenol, and the acid addition salts thereof.

[037] Examples of ortho-aminophenols that may be mentioned, are 2-aminophenol, 2-amino-5-methylphenol, 2-amino-6-methylphenol, 5-acetamido-2-aminophenol, and the acid addition salts thereof.

[038] Examples of heterocyclic bases that may be mentioned, are pyridine derivatives, pyrimidine derivatives and pyrazole derivatives.

[039] Examples of pyridine derivatives that may be mentioned are the compounds described, in patents GB 1 026 978 and GB 1 153 196, such as 2,5-diaminopyridine; as well as 2-(4-methoxyphenyl)amino-3-aminopyridine, 2,3-diamino-6-methoxypyridine,

2-(β -methoxyethyl)amino-3-amino-6-methoxypyridine and 3,4-diaminopyridine, and the acid addition salts thereof.

[040] Other pyridine oxidation bases that may be useful in the dye composition disclosed herein are 3-aminopyrazolo[1,5-a]pyridine oxidation bases and the addition salts thereof described, for example, in patent application FR 2 801 308. By way of example, mention may be made of pyrazolo[1,5-a]pyrid-3-ylamine;

2-acetylaminopyrazolo[1,5-a]pyrid-3-ylamine; 2-morpholin-4-ylpyrazolo[1,5-a]pyrid-3-ylamine; 3-aminopyrazolo[1,5-a]pyridine-2-carboxylic acid; 2-methoxypyrazolo[1,5-a]pyrid-3-ylamine; (3-aminopyrazolo[1,5-a]pyrid-7-yl)methanol; 2-(3-aminopyrazolo[1,5-a]pyrid-5-yl)ethanol; 2-(3-aminopyrazolo[1,5-a]pyrid-7-yl)ethanol; (3-aminopyrazolo[1,5-a]pyrid-2-yl)-methanol; 3,6-diaminopyrazolo[1,5-a]pyridine; 3,4-diaminopyrazolo[1,5-a]pyridine; pyrazolo[1,5-a]pyridine-3,7-diamine; 7-morpholin-4-ylpyrazolo[1,5-a]pyrid-3-ylamine; pyrazolo[1,5-a]pyridine-3,5-diamine; 5-morpholin-4-ylpyrazolo[1,5-a]pyrid-3-ylamine; 2-[(3-aminopyrazolo[1,5-a]pyrid-5-yl)(2-hydroxyethyl)amino]ethanol; 2-[(3-aminopyrazolo[1,5-a]pyrid-7-yl)(2-hydroxyethyl)amino]ethanol; 3-aminopyrazolo[1,5-a]pyrid-5-ol; 3-aminopyrazolo[1,5-a]pyrid-4-ol; 3-aminopyrazolo[1,5-a]pyrid-6-ol; 3-aminopyrazolo[1,5-a]pyrid-7-ol; and the addition salts thereof with an acid or with a base.

[041] Additional examples of pyrimidine derivatives that may be mentioned are the compounds described, for instance, in patents DE 2 359 399; JP 88-169 571; JP 05-63124; EP 0 770 375 or patent application WO 96/15765, such as 2,4,5,6-tetraaminopyrimidine, 4-hydroxy-2,5,6-triaminopyrimidine, 2-hydroxy-4,5,6-triaminopyrimidine, 2,4-dihydroxy-5,6-diaminopyrimidine, and 2,5,6-triaminopyrimidine. Further mention may be made of pyrazolopyrimidine derivatives, such as those mentioned in patent application FR-A-2 750 048, for example, pyrazolo[1,5-a]pyrimidine-3,7-diamine; 2,5-

dimethylpyrazolo[1,5-a]pyrimidine-3,7-diamine; pyrazolo[1,5-a]pyrimidine-3,5-diamine; 2,7-dimethylpyrazolo[1,5-a]pyrimidine-3,5-diamine; 3-aminopyrazolo[1,5-a]pyrimidin-7-ol; 3-aminopyrazolo[1,5-a]pyrimidin-5-ol; 2-(3-aminopyrazolo[1,5-a]pyrimidin-7-ylamino)ethanol, 2-(7-aminopyrazolo[1,5-a]pyrimidin-3-ylamino)ethanol, 2-[(3-aminopyrazolo[1,5-a]pyrimidin-7-yl)(2-hydroxyethyl)amino]ethanol, 2-[(7-aminopyrazolo[1,5-a]pyrimidin-3-yl)(2-hydroxyethyl)amino]ethanol, 5,6-dimethylpyrazolo[1,5-a]pyrimidine-3,7-diamine, 2,6-dimethylpyrazolo[1,5-a]pyrimidine-3,7-diamine, 2,5,N7,N7-tetramethylpyrazolo[1,5-a]pyrimidine-3,7-diamine and 3-amino-5-methyl-7-imidazolylpropylaminopyrazolo[1,5-a]pyrimidine, the acid addition salts thereof, and the tautomeric forms thereof, when a tautomeric equilibrium exists.

[042] Examples of pyrazole derivatives that may be mentioned are the compounds described in patents DE 3 843 892 and DE 4 133 957 and patent applications WO 94/08969, WO 94/08970, FR-A-2 733 749 and DE 195 43 988, for instance, 4,5-diamino-1-methylpyrazole, 4,5-diamino-1-(β -hydroxyethyl)pyrazole, 3,4-diaminopyrazole, 4,5-diamino-1-(4'-chlorobenzyl)pyrazole, 4,5-diamino-1,3-dimethylpyrazole, 4,5-diamino-3-methyl-1-phenylpyrazole, 4,5-diamino-1-methyl-3-phenylpyrazole, 4-amino-1,3-dimethyl-5-hydrazinopyrazole, 1-benzyl-4,5-diamino-3-methylpyrazole, 4,5-diamino-3-tert-butyl-1-methylpyrazole, 4,5-diamino-1-tert-butyl-3-methylpyrazole, 4,5-diamino-1-(β -hydroxyethyl)-3-methylpyrazole, 4,5-diamino-1-ethyl-3-methylpyrazole, 4,5-diamino-1-ethyl-3-(4'-methoxyphenyl)pyrazole, 4,5-diamino-1-ethyl-3-hydroxymethylpyrazole, 4,5-diamino-3-hydroxymethyl-1-methylpyrazole, 4,5-diamino-3-hydroxymethyl-1-isopropylpyrazole, 4,5-diamino-3-methyl-1-isopropylpyrazole, 4-amino-5-(2'-aminoethyl)amino-1,3-dimethylpyrazole, 3,4,5-triaminopyrazole, 1-methyl-3,4,5-triaminopyrazole, 3,5-diamino-1-

methyl-4-methylaminopyrazole, 3,5-diamino-4-(β -hydroxyethyl)amino-1-methylpyrazole, and the acid addition salts thereof.

[043] The optional at least one additional oxidation base present in the dye composition disclosed herein can be present in the composition in an amount ranging approximately from 0.001% to 10%, by weight relative, for each additional oxidation base, to the total weight of the dye composition. For example, the at least one additional oxidation base may be present in the dye composition in an amount ranging approximately from 0.005% to 6% by weight relative, for each additional oxidation base, to the total weight of the dye composition.

[044] In general, the addition salts of the oxidation bases and of the couplers that may be used in the dye composition disclosed herein are chosen from the acid addition salts, such as the hydrochlorides, hydrobromides, sulphates, citrates, succinates, tartrates, lactates, tosylates, benzenesulphonates, phosphates and acetates, and the base addition salts, such as sodium hydroxide, potassium hydroxide, ammonia, amines and alkanolamines.

[045] In another embodiment, the dye composition may also comprise at least one direct dye that may be chosen from nitrobenzene dyes, azo direct dyes and methine direct dyes. The direct dyes may be of nonionic, anionic or cationic nature.

[046] A medium that is suitable for dyeing, also known as the dye support, generally comprises water or a mixture of water and at least one organic solvent to dissolve the compounds which would not be sufficiently soluble in water. Examples of organic solvents that may be used are C₁-C₄ lower alkanols, (such as ethanol and isopropanol); polyols and polyol ethers (such as 2-butoxyethanol, propylene glycol, propylene glycol

monomethyl ether, diethylene glycol monoethyl ether and monomethyl ether), and aromatic alcohols (such as benzyl alcohol or phenoxyethanol), and mixtures thereof.

[047] The organic solvents may be present in the dye composition in proportions ranging approximately from 1% to 40% by weight, relative to the total weight of the dye composition; for example, ranging approximately from 5% to 30% by weight, relative to the total weight of the dye composition.

[048] The dye composition as disclosed herein can also contain at least one of various adjuvants conventionally used in compositions for dyeing the hair, such as anionic, cationic, nonionic, amphoteric and zwitterionic surfactants, and mixtures thereof; anionic, cationic, nonionic, amphoteric and zwitterionic polymers, and mixtures thereof; inorganic and organic thickeners, for example, anionic, cationic, nonionic and amphoteric associative polymeric thickeners; antioxidants; penetration agents; sequestering agents; fragrances; buffers; dispersing agents; packaging agents such as, for example, silicones, which may or may not be volatile or modified; film-forming agents; ceramides; preserving agents and opacifiers.

[049] The at least one adjuvant may be present in the dye composition in an amount ranging approximately from 0.01% to 20% by weight relative, for each adjuvant, to the weight of the composition.


[050] Needless to say, a person skilled in the art will take care to select any of the optional additional compounds described above such that the advantageous properties intrinsically associated with the oxidation dye as disclosed herein are not, or are not substantially, adversely affected by the addition(s) envisaged.

[051] The pH of the dye composition as disclosed herein, may range approximately from 3 to 12 and, for example, can range approximately from 5 to 11. It may

be adjusted to the desired value using acidifying or basifying agents usually used in the dyeing of keratin fibers, or alternatively using standard buffer systems.

[052] Among the acidifying agents that may be mentioned, for example, are inorganic and organic acids such as hydrochloric acid; orthophosphoric acid; sulphuric acid; carboxylic acids such as acetic acid tartaric acid, citric acid and lactic acid; and sulphonic acids.

[053] Among the basifying agents that may be mentioned, for example, are aqueous ammonia, alkaline carbonates, alkanolamines such as mono-, di- and triethanolamine and derivatives thereof, sodium hydroxide, potassium hydroxide and the compounds of formula (II) below:



wherein W is a propylene residue that can be unsubstituted or substituted with an entity chosen from hydroxyl groups and C₁-C₄ alkyl radicals; R_a, R_b, R_c and R_d, which may be identical or different, are each chosen from hydrogen atoms, C₁-C₄ alkyl radicals, and C₁-C₄ hydroxyalkyl radicals.

[054] The dye composition as disclosed herein may be in various forms, such as liquids, creams and gels, or in any other form that is suitable for dyeing keratin fibers, for example human hair.

[055] Also disclosed herein is a process for dyeing keratin fibers by of applying the dye composition as defined above to keratin fibers, in the presence of an oxidizing agent for a time that is sufficient to develop the desired coloration. The color may be developed at acidic, neutral and alkaline pH. The oxidizing agent may be mixed with the composition of

the invention just at the time of use, or it may be used starting with an oxidizing composition comprising it, which can be applied simultaneously or sequentially to the dye composition of the invention.

[056] According to one embodiment, the dye composition as described above is mixed, for example, at the time of use, with a composition comprising, in a medium that is suitable for dyeing, at least one oxidizing agent, this oxidizing agent being present in an amount that is sufficient to develop a coloration. The mixture obtained is then applied to the keratin fibers. After an action time ranging approximately from 3 to 50 minutes, for example, ranging approximately from 5 to 30 minutes, the keratin fibers are rinsed, washed with shampoo, rinsed again and then dried.

[057] The oxidizing agents conventionally used for the oxidation dyeing of keratin fibers are, for example, hydrogen peroxide, urea peroxide, alkali metal bromates, persalts (such as perborates and persulphates), peracids and oxidase enzymes, among which mention may be made of peroxidases, 2-electron oxidoreductases such as uricases, and 4-electron oxygenases, for instance laccases. In one embodiment of the oxidizing composition disclosed herein, hydrogen peroxide is the oxidizing agent.

[058] The oxidizing composition may also contain at least one adjuvant conventionally used in compositions for dyeing the hair and as defined above.


[059] The pH of the oxidizing composition comprising the oxidizing agent is such that, after mixing with the dye composition, the pH of the resulting composition applied to the keratin fibers ranges approximately from 3 to 12, for example; the pH can range approximately from 5 to 11. It may be adjusted to the desired value by means of acidifying or basifying agents usually used in the dyeing of keratin fibers and as defined above.

[060] The ready-to-use dye composition that is finally applied to the keratin fibers may be in various forms, such as in the form of liquids, creams or gels or any other form that is suitable for dyeing keratin fibers, for instance, human hair.

[061] Another embodiment disclosed herein is a multi-compartment dyeing device or "kit", in which a first compartment contains the dye composition of the present invention defined above and a second compartment contains an oxidizing agent. This device may be equipped with a means for applying the desired mixture to the hair, such as the devices described in patent FR-2 586 913, assigned to L'Oréal.

[062] Using this device, it is possible to dye keratin fibers using a process that involves mixing a dye composition comprising at least one oxidation base comprising at least one pyrrolidyl -substituted para-phenylenediamine derivative chosen from derivatives corresponding to formula (I), with an oxidizing agent, and applying the mixture obtained to the keratin fibers for a time that is sufficient to develop the desired coloration.

[063] The examples that follow serve to illustrate the invention without, however, being limiting in nature.

EXAMPLES**SYNTHESIS EXAMPLES****Example 1:** Synthesis of 4-(3-imidazolyl-1-ylpyrrolid-1-yl)phenylamine**I. Synthesis of 1-(4-nitrophenyl)pyrrolid-3-yl methanesulphonate (2)**

[064] 40 ml (0.516 mol) of mesyl chloride were added dropwise at 5°C to 83.3 g (0.4 mol) of N-(4-nitrophenyl)-3-hydroxypyrrolidine dissolved in 625 ml of anhydrous THF and 72.7 ml (0.6 mol) of triethylamine. The mixture was allowed to return to room temperature and was then poured into ice to form a precipitate.

[065] After suction-filtration and drying of the precipitate, 109 g of yellow powder

[066] (2) were obtained.

Melting point = 203°C

¹H NMR (400 MHz-DMSO) ppm 8.09 (d, 2H); 6.68 (d, 2H); 5.47 (m, 1H); 3.77-3.48 (m, 4H); 3.28 (s, 3H); 2.35 (m, 2H).

II. Synthesis of 1-[1-(4-nitrophenyl)pyrrolid-3-yl]-1H-imidazole (3)

[067] 22 g (0.0767 mol) of 1-(4-nitrophenyl)pyrrolid-3-yl methanesulphonate (2) were heated at 95°C for 2 hours in 170 g of imidazole (2.5 mol). This mixture was poured into 1 l of ice-cold water until crystallization takes place. After filtration and drying, the yellow powder obtained was chromatographed, eluting with dichloromethane, and 14.5 g of derivative (3) were recovered (73.2% yield).

Melting point = 163°C

¹H NMR (400 MHz-DMSO) ppm 8.20 (m, 2H); 7.87 (s, 1H); 7.38 (s, 1H); 7.06 (s, 1H); 6.81 (m, 2H); 5.17 (m, 1H); 4.06 (m, 1H); from 3.79 to 3.65 (m, 3H); 2.66 (m, 1H); 2.50 (m, 1H);


Mass ESI+: m/z = 259 [M+]

III. Synthesis of 4-(3-imidazolyl-1-ylpyrrolid-1-yl)phenylamine hydrochloride (4)

[068] 13.5 g (0.0522 mol) of the above derivative dissolved in 700 ml of ethanol were hydrogenated in the presence of palladium on charcoal under a hydrogen pressure of 8 bar. After filtering off the catalyst, the expected derivative (4) was isolated in the form of the hydrochloride. 13 g of white powder are obtained; 82.7% yield.

¹H NMR (400 MHz-DMSO) ppm 9.29 (s, 1H); 7.84 (t, 1H); 7.71 (t, 1H); 7.26 (m, 2H); 6.67 (m, 2H); 5.29 (m, 1H); 3.77 (m, 1H); 3.68 (m, 2H); 3.38 (m, 1H); 2.51 (m, 1H); 2.46 (m, 1H);

Mass ESI+: m/z = 229 [M+]

Example 2: Synthesis of 4-([1,3']bipyrrolinyl-1'-yl)phenylamine**I. Synthesis of 1'-(4-nitrophenyl)[1,3]bipyrrolidinyl (5)**

[069] 5 g (0.0174 mol) of 1-(4-nitrophenyl)pyrrolidin-3-yl methanesulphonate (2) were heated at 85°C for 2 hours in 30 ml of pyrrolidine (mol). This mixture was poured into ice-cold water until crystallization takes place. After filtration and drying, the yellow powder obtained was chromatographed, eluting with dichloromethane/methanol (98/2), and 2.6 g of derivative (5) were recovered (53% yield).

Melting point = 114°C

¹H NMR (400 MHz-DMSO) ppm 8.04 (m, 2H); 6.61 (m, 2H); 3.60 (m, 2H); 3.40 (m, 1H); 3.24 (m, 1H); 2.86 (m, 1H); 2.50 (m, 2H); 2.16 (m, 1H); 1.92 (m, 1H); 1.70 (m, 4H).


Mass ESI+: m/z = 262 [M+]

II. Synthesis of 4-([1,3']bipyrrolinyl-1'-yl)phenylamine hydrochloride (6)

[070] 2.5 g (0.0096 mol) of the above derivative (5) dissolved in 400 ml of ethanol were hydrogenated in the presence of palladium on charcoal under a hydrogen pressure of 8 bar at a temperature of 50°C; after filtering off the catalyst, the expected derivative (6) was isolated in the form of the hydrochloride. 1.3 g of a white powder are obtained; 44% yield.

¹H NMR (400 MHz-D₂O) ppm 7.33 (m, 2H); 6.86 (m, 2H); 4.11 (m, 1H); 3.75 (m, 3H); 3.60 (m, 2H); 3.39 (m, 1H); 2.60 (m, 1H); 2.31 (m, 1H); 2.18 (m, 2H); 2.06 (m, 2H).

Mass ESI+: m/z = 232 [M+]

Example 3: Synthesis of [1-(4-aminophenyl)pyrrolid-3-yl](3-imidazol-1-ylpropyl)amine

I. Synthesis of [3-(imidazol-1-yl)propyl][1-(4-nitrophenyl)pyrrolid-3-yl]amine (7)

[071] 30 g (0.105 mol) of 1-(4-nitrophenyl)pyrrolid-3-yl methylsulphonate (**2**) were heated at 90°C for 18 hours with 120 g of aminopropylimidazole (0.958 mol). This mixture was poured into ice-cold water and the product was extracted with dichloromethane. The yellow powder obtained was chromatographed, eluting with dichloromethane/methanol (98/2), and 15.2 g of derivative (**7**) were recovered (48.2% yield).

Melting point = 74°C

¹H NMR (400 MHz-DMSO) ppm 8.06 (m, 2H); 7.60 (s, 1H); 7.15 (s, 1H); 6.84 (s, 1H); 6.61 (m, 2H); 4.35 (bs, 1H); 4.03 (m, 2H); 3. (m, 2H); 3.49 (m, 2H); 3.24 (m, 1H); 2.15 (m, 1H); 1.85 (m, 3H).

Mass ESI⁺: m/z = 316 [M⁺]

II. Synthesis of [1-(4-aminophenyl)pyrrolid-3-yl](3-imidazol-1-ylpropyl)amine hydrochloride (8)

[072] 6.5 g (0.0206 mol) of the above derivative (**7**) dissolved in 500 ml of ethanol were hydrogenated in the presence of palladium on charcoal, under 10 bar; after filtering off the catalyst, the expected derivative (**8**) was isolated in the form of the hydrochloride.

7.07 g of white powder are obtained; 86% yield.

¹H NMR (400 MHz-D₂O) ppm 8.86 (s, 1H); 7.63 (d, 1H); 7.55 (d, 1H); 7.34 (d, 2H); 6.83 (d, 2H); 4.46 (t, 1H); 4.13 (m, 1H); 3.72 (m, 1H); 3.61 (m, 2H); 3.40 (m, 1H); 2.58 (m, 1H); 2.43 (m, 1H); 2.31 (m, 1H).

Mass ESI⁺: m/z = 286 [M⁺]

EXAMPLES OF DYEING**EXAMPLES 1 TO 11: DYEING IN ALKALINE MEDIUM**

Examples	1	2	3	4	5	6	7	8	9	10	11
4-(3-Imidazol-1-yl-pyrrolid-1-yl)phenylamine (base)	10 ⁻³	10 ⁻³	10 ⁻³	10 ⁻³	-	-	-	-	-	-	-
[1-(4-Aminophenyl)pyrrolid-3-yl]-(3-imidazol-1-ylpropyl)amine (base)	-	-	-	-	10 ⁻³	10 ⁻³	10 ⁻³	-	-	-	-
4-[1,3']Bipyrrolid-1'-ylphenylamine (base)	-	-	-	-	-	-	-	10 ⁻³	10 ⁻³	10 ⁻³	10 ⁻³
2-(2,4-Diaminophenoxy)ethanol dihydrochloride (coupler)	10 ⁻³	-	-	-	10 ⁻³	-	-	10 ⁻³	-	-	-
3-Amino-2-chloro-6-methylphenol hydrochloride (coupler)	-	10 ⁻³	-	-	-	-	-	-	10 ⁻³	-	-
3,6-Dimethyl-1H-pyrazolo[5,1-c][1,2,4]triazole (coupler)	-	-	10 ⁻³	-	-	10 ⁻³	-	-	-	10 ⁻³	-
2-methyl-5-aminophenol (coupler)	-	-	-	10 ⁻³	-	-	10 ⁻³	-	-	-	10 ⁻³
Dye support (1)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Demineralized water qs	100 g	100 g	100 g	100 g	100 g	100 g	100 g	100 g	100 g	100 g	100 g

- the amounts of base and of coupler are expressed in moles

(*) Dye support (1) pH 9.5

96° ethyl alcohol	20.8	g
Sodium metabisulphite as an aqueous 35% solution	0.23	g A.M.
Pentasodium salt of diethylenetriaminepenta-acetic acid, as an aqueous 40% solution	0.48	g A.M.
C ₈ -C ₁₀ alkyl polyglucoside as an aqueous 60% solution	3.6	g A.M.
Benzyl alcohol	2.0	g
Polyethylene glycol containing 8 units of ethylene oxide	3.0	g
NH ₄ Cl	4.32	g
Aqueous ammonia containing 20% NH ₃	2.94	g

[073] At the time of use, each composition was mixed with an equal weight of 20-volumes aqueous hydrogen peroxide solution (6% by weight). A final pH of 9.5 was obtained.

[074] Each mixture obtained was applied to locks of grey hair containing 90% white hairs. After an action time of 30 minutes, the locks were rinsed, washed with a standard shampoo, rinsed again and then dried.

[075] The dyeing results below were obtained.

Examples	1	2	3	4	5	6
Shade observed	Violet-blue	Blue-violet	Red-violet	Violet	Violet-blue	Red-violet

Examples	7	8	9	10	11
Shade observed	Violet	Violet-blue	Blue-violet	Chromatic red-violet	Violet

EXAMPLES 12 TO 29: DYEING IN ACIDIC MEDIUM

The dye compositions below were prepared:

Examples	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29
4-(3-Imidazol-1-ylpyrrolid-1-yl)phenylamine (base)	10 ⁻³	10 ⁻³	10 ⁻³	10 ⁻³	10 ⁻³	10 ⁻³	-	-	-	-	-	-	-	-	-	-	-	-
[1-(4-Aminophenyl)pyrrolid-3-yl]-(3-imidazol-1-ylpropyl)-amine (base)	-	-	-	-	-	-	10 ⁻³	10 ⁻³	10 ⁻³	10 ⁻³	10 ⁻³	10 ⁻³	-	-	-	-	-	-
4-[1,3']Bipyrrlid-1'-ylphenylamine (base)	-	-	-	-	-	-	-	-	-	-	-	-	10 ⁻³	10 ⁻³	10 ⁻³	10 ⁻³	10 ⁻³	10 ⁻³
2-(2,4-Diaminophenoxy)ethanol dihydrochloride (coupler)	10 ⁻³	-	-	-	-	-	10 ⁻³	-	-	-	-	-	10 ⁻³	-	-	-	-	-
3-Amino-2-chloro-6-methylphenol hydrochloride (coupler)	-	10 ⁻³	-	-	-	-	-	10 ⁻³	-	-	-	-	-	10 ⁻³	-	-	-	-
2-Methyl-5-aminophenol (coupler)	-	-	10 ⁻³	-	-	-	-	-	10 ⁻³	-	-	-	-	-	10 ⁻³	-	-	-
2-Amino-3-pyridinol (coupler)	-	-	-	10 ⁻³	-	-	-	-	-	10 ⁻³	-	-	-	-	-	10 ⁻³	-	-
3,6-Dimethyl-1H-pyrazolo [5,1-c][1,2,4]triazole (coupler)	-	-	-	-	10 ⁻³	-	-	-	-	-	10 ⁻³	-	-	-	-	-	10 ⁻³	-
6-Hydroxy-1H-indole (coupler)	-	-	-	-	-	10 ⁻³	-	-	-	-	-	10 ⁻³	-	-	-	-	-	10 ⁻³
Dye support (2)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)	(*)
Deminerlized water qs	100g	100g	100g	100g	100g	100g	100g	100g	100g	100g	100g	100g	100g	100g	100g	100g	100g	100g

- The amounts of base and of coupler are expressed in moles

(*) Dye support (2) pH 7

96° ethyl alcohol	20.0	g
Sodium metabisulphite as an aqueous 35% solution	0.23	g A.M.
Pentasodium salt of diethylenetriaminepenta-acetic acid as an aqueous 40% solution	0.48	g A.M.
C ₈ -C ₁₀ alkyl polyglucoside as an aqueous 60% solution	3.6	g A.M.
Benzyl alcohol	2.0	g
Polyethylene glycol containing 8 units of ethylene oxide	3.0	g
Na ₂ HPO ₄	0.28	g
KH ₂ PO ₄	0.46	g

[076] At the time of use, each composition was mixed with an equal weight of 20-volumes aqueous hydrogen peroxide solution (6% by weight). A final pH of 7 was obtained.

[077] Each mixture obtained was applied to locks of grey hair containing 90% white hairs. After an action time of 30 minutes, the locks were rinsed, washed with a standard shampoo, rinsed again and then dried.

[078] The following dyeing results were obtained.

Examples	12	13	14	15	16	17
Shade observed	Violet-blue	Blue-violet	Violet	Grey-violet	Red-violet	Grey-violet

Examples	18	19	20	21	22	23
Shade observed	Violet-blue	Blue-violet	Violet	Grey-violet	Red-violet	Grey-violet

Examples	24	25	26	27	28	29
Shade observed	Violet-blue	Blue-violet	Violet	Grey-violet	Red-violet	Grey-violet